



## 2. Andrejewski Vorlesung/ SFB Seminartag

### ZEIT:

10.1.2006, 14:00 Uhr - 19:00 Uhr

### ORT:

im AEI in Golm, Max-Planck-Campus  
Am Mühlenberg 1, 14476 Golm  
Central Building, room number Z-050

### PROGRAMM:

14:00                    **Prof. Dr. Albrecht Klemm, University of Wisconsin,  
Madison**

#### **Mirror symmetry and the topological A- and B-model**

Mirror symmetry on CY manifolds exchanges the symplectic structure on  $M$ , actually a complexified Kähler structure, with the complex structure on a mirror dual CY manifold  $W$ . The deformation theory of each of these structures can be described by a topological string theory called the topological A- and the B-model respectively. These models are cohomological theories defined by nilpotent operators  $Q_A$  and  $Q_B$ . We will show that  $Q_A$  exists on every symplectic manifold while  $Q_B$  exists only on CY manifolds and certain generalizations thereof. The latter fact is related to the Tian & Todorov theorem on the unobstructedness of complex structure deformations on CY spaces and generalizations by Hitchin. We will then discuss properties of cohomological theories notably the descend- and topological recursion relations. The solution of the topological B-model using these recursion relations and some classical methods of complex structure deformation theory on  $W$  is worked out and related by mirror symmetry to the Gromov-Witten theory captured by the topological A-model on  $M$ .

16:00                    **Prof. Dr. Jens Hoppe (Royal Institute of Technology,  
Stockholm), guest of the AEI**

#### **Kontakt:**

Humboldt-Universität zu Berlin . Institut für Mathematik  
SFB 647 . Unter den Linden 6 . 10099 Berlin  
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727  
sfb647@math.hu-berlin.de

[www.raumzeitmaterie.de](http://www.raumzeitmaterie.de)

## Aspects of Membrane Dynamics

17:30

**Dr. Simon Chiossi (HU)**

### **G2 structures on solvmanifolds**

Conformally G2 manifolds are Riemannian manifolds with a G2 structure whose metric can be modified to a holonomy structure by a conformal change. There is an interesting construction of homogeneous conformally G2 structures on solvmanifolds built from underlying SU(3) structures. I will show how the corresponding non-homogeneous G2 metrics can be obtained also by evolving the SU(3) structure in time. (Possible reference: arXive math.DG/0510087)

#### **Kontakt:**

Humboldt-Universität zu Berlin . Institut für Mathematik  
SFB 647 . Unter den Linden 6 . 10099 Berlin  
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727  
sfb647@math.hu-berlin.de

[www.raumzeitmaterie.de](http://www.raumzeitmaterie.de)