



Dr. Norbert Peyerimhoff Pompeiu's Problem on Damek-Ricci spaces

ZEIT:

7.11.2007, 16:30 Uhr

ORT:

HU-Berlin
Institut für Mathematik
Rudower Chaussee 25
Raum 1.410

In 1929, the Rumanian mathematician Dimitrie Pompeiu asked the following question: given a continuous function f on \mathbb{R}^2 and a compact set K . Assume that the integral of f over all images of K under rigid motions vanishes. Does this imply that the function f itself is zero? The answer is negative in the case that K is a disk. But it can be shown that the conclusion holds in the case that f vanishes on all disks of radius r_1 and of radius r_2 , as long as r_1 and r_2 avoid a certain countable set of radii. It is natural to ask similar questions in more general geometries. In this talk we discuss the same (two radius) problem in Damek-Ricci spaces. These spaces became famous as counterexamples of the Lichnerowitch conjecture, namely that every harmonic space should be a rank one symmetric space.

Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de

www.raumzeitmaterie.de