

Dr. Anne-Sophie Kaloghiros

Non-factorial Fano 3-folds

TIME:

26 May 2008, 16:15

LOCATION:

Freie Universität Berlin
Institut für Mathematik
Arnimallee 3, Rm. 119
114195 Berlin

A singular variety is \mathbb{Q} -factorial if a multiple of every Weil divisor (a codimension 1 cycle) is a Cartier divisor (i.e. can be defined locally by one equation). \mathbb{Q} -factoriality is a subtle topological property that depends both on the local analytic type of singularities and on their relative position. The purpose of these talks will be to study non-factorial Fano 3-folds with mild singularities. Fano 3-folds are the higher dimensional analogues of spheres, they have 'positive curvature', and as such play an important role in the Minimal Model program. I will show that one can understand to some extent how far Fano 3-folds with terminal Gorenstein singularities can be from being factorial. In the first talk, I will first give an introduction to terminal/canonical/Gorenstein singularities, and to factoriality. I will also present the Minimal Model Program for (mildly singular) 3-folds. In the second talk, I will apply these techniques to the study of non-factorial Fano 3-folds.

Contact:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de