



Burglind Juhl-Jöricke

Analytic knots, satellites and the 4-ball genus

TIME:

6 Jul 2012, 10:00 - 11:00

LOCATION:

HU

Institut für Mathematik, Raum 1.114

Rudower Chaussee 25

12489 Berlin

Abstract. A satellite of a knot K is a link contained in a tubular neighbourhood of K (but not in a 3-ball contained in this neighbourhood), and not isotopic to K . In a classical paper Schubert gives a lower bound for the genus of satellites.

Consider a knot or link in the unit sphere in \mathbb{C}^2 . Call it analytic (respectively, smoothly analytic) if it bounds a complex curve (respectively, a smooth complex curve) in the complex ball.

Let K be a smoothly analytic knot. For analytic satellite links contained in a sufficiently small tubular neighbourhood of K there is a (sharp) lower bound of the 4-ball genus (but not of the genus).

Moreover, these links can be completely described. The problem is related to branched coverings and braided surfaces.

Contact:

Humboldt-Universität zu Berlin . Institut für Mathematik

SFB 647 . Unter den Linden 6 . 10099 Berlin

Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727

sfb647@math.hu-berlin.de

www.raumzeitmaterie.de